



Aminoquinolines afford resistance to cerebral malaria in susceptible mice[☆]



Jelena Srbljanović^a, Branko Bobić^a, Tijana Štajner^a, Aleksandra Uzelac^a, Igor Opsenica^b, Nataša Terzić-Jovanović^c, Neda Bauman^a, Bogdan A. Šolaja^{b,d}, Olgica Djurković-Djaković^{a,*}

^a Institute for Medical Research, University of Belgrade, Dr. Subotića 4, P.O. Box 39, 11129 Belgrade, Serbia

^b Faculty of Chemistry, University of Belgrade, Studentski trg 16, P.O. Box 51, 11158 Belgrade, Serbia

^c Institute of Chemistry, Technology and Metallurgy, 11000 Belgrade, Serbia

^d Serbian Academy of Sciences and Arts, Knez Mihailova 35, 11000 Belgrade, Serbia

ARTICLE INFO

Article history:

Received 16 January 2020

Received in revised form 7 July 2020

Accepted 31 July 2020

Available online 15 August 2020

Keywords:

Malaria
Aminoquinolines
Plasmodium berghei
C57BL/6 mice
Hyperparasitaemia

ABSTRACT

Objectives: Malaria treatment is impeded by increasing resistance to conventional antimalarial drugs. Here we explored the activity of ten novel benzothiofene, thiophene and benzene aminoquinolines. **Methods:** In vitro testing was performed by the lactate dehydrogenase assay in chloroquine (CQ)-sensitive *Plasmodium falciparum* strain 3D7 and CQ-resistant (CQ^R) *P. falciparum* strain Dd2. In vivo activity was evaluated by a modified Thompson test using C57BL/6 mice infected with *Plasmodium berghei* ANKA strain.

Results: Nine of the ten compounds had a lower 50% inhibitory concentration (IC₅₀) than CQ against the CQ^R strain Dd2. Five of these compounds that were available for in vivo evaluation were shown to be non-toxic. All five compounds administered at a dose of 160 mg/kg/day for 3 days prolonged the survival of treated compared with untreated mice. Untreated control mice died by Day 7 with a mean parasitaemia of 15%. Among treated mice, a dichotomous outcome was observed, with a two-third majority of treated mice dying by Day 17 with a low mean parasitaemia of 5%, whilst one-third survived longer with a mean hyperparasitaemia of 70%; specifically, five of these mice survived a mean of 25 days, whilst two even survived past Day 31.

Conclusions: The significant antimalarial potential of this aminoquinoline series is illustrated by its excellent in vitro activity against the CQ^R *P. falciparum* strain and significant in vivo activity. Interestingly, compounds CIAQ7, CIAQ9 and CIAQ11 were able to confer resistance to cerebral malaria and afford a switch to hyperparasitaemia to mice prone to the neurological syndrome.

© 2020 The Author(s). Published by Elsevier Ltd on behalf of International Society for Antimicrobial Chemotherapy. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Malaria is a parasitic infection caused by protozoa of the genus *Plasmodium* that affects millions of people worldwide [1]. The resistance both of parasites and vectors to existing drugs is an ongoing and increasing problem, hampering both prevention and treatment of malaria. Indeed, *Plasmodium* parasites are developing resistance to nearly all conventional antimalarials, and *Anopheles* vectors are becoming resistant to insecticides, with

no vaccine existing to date. Additionally, climate changes and migrations may favour further spread and the re-emergence of malaria even in areas considered free of the disease. Given these circumstances, discovering new antimalarial compounds that can overcome parasite resistance has emerged as a global scientific goal [2–7].

Our recent work focused on the antimalarial effect of 26 novel aminoquinolines investigated in both in vitro and in vivo model systems and showed a great potential of adamantane derivatives, with two compounds shown to cure *Plasmodium berghei*-infected mice [8]. The current study examined ten novel benzothiofene, thiophene and benzene derivatives of the same aminoquinoline series, whose synthesis has recently been described [9–11]. Here we report on the activity of these ten aminoquinolines in different in vitro model systems as well as in vivo and note that some

[☆] The results of this study were presented in part at the 14th International Congress of Parasitology (ICOPA2018), 19–24 August 2018, Daegu, South Korea.

* Corresponding author.

E-mail address: olgicadj@imi.bg.ac.rs (O. Djurković-Djaković).

conferred resistance to cerebral malaria and a switch to hyperparasitaemia of mice prone to the neurological syndrome.

2. Materials and methods

2.1. Parasites

For in vitro drug assays, chloroquine (CQ)-sensitive (CQ^S) *Plasmodium falciparum* strain 3D7 and CQ-resistant (CQ^R) *P. falciparum* strain Dd2 were used. Cell cultures were maintained as described previously [12]. Parasites were synchronised with 5% sorbitol and ring-stage parasites were seeded in 96-well plates to achieve 2% parasitaemia and 0.75% haematocrit.

For in vivo testing, *P. berghei* ANKA strain maintained through serial intraperitoneal (i.p.) passage in C57BL/6 mice was used.

2.2. Mice

Female C57BL/6 mice (Medical Military Academy Animal Research Facility, Belgrade, Serbia) weighing 19–21 g were used. Mice were housed in groups of four to six animals in the Institute for Medical Research Animal Facility under a natural photoperiod and were offered drinking water and standard feed ad libitum.

2.3. Compounds

Ten novel aminoquinoline derivatives with chemical modifications at the aminoquinoline moiety and side chain, synthesised at the Faculty of Chemistry, University of Belgrade [9–11], were examined (Table 1). According to the structure of the aminoquinoline moiety, compounds were classified into 4-aminoquinolines (AQs) (two compounds) and 7-chloro-4-aminoquinolines (CIAQs) (eight compounds).

For experimental use in vitro, compounds were dissolved in dimethyl sulfoxide (DMSO) at a stock concentration of 50 mM. Compounds were further diluted in complete RPMI 1640 culture medium and the final DMSO concentration was $\leq 0.2\%$.

For in vivo experiments, compounds were suspended in 0.5% hydroxyethyl cellulose/0.1% Tween 80 and were administered orally (p.o.).

2.4. Experimental design

All compounds were screened in vitro and, if appropriate, were titrated to obtain 50% inhibitory concentration (IC₅₀) values and examined for antimalarial efficacy in vivo. Prior to in vivo evaluation of the selected compounds, all were tested for toxicity in mice.

2.5. Assays

In vitro evaluation of the antimalarial activity of the compounds was performed using the colorimetric lactate dehydrogenase assay [13]. Compounds were first screened at a concentration of 500 nM (three replicates per compound for each *P. falciparum* strain) and those that showed a minimum of 50% parasite growth inhibition were further titrated to determine their IC₅₀ value. CQ was used as a positive control. Briefly, compounds were titrated using eight two-fold dilutions, starting at a concentration of 256 nM or 1000 nM, depending on the results of the initial screening. Three independent experiments were performed for each compound, with three replicates per compound for each *P. falciparum* strain. More precisely, the IC₅₀ value of each compound and CQ was calculated as the geometric mean IC₅₀ value of the individual experiment for a particular strain of *P. falciparum*.

In vivo toxicity of the experimental compounds was examined through a series of experiments in which selected compounds were administered at a dose of 160 mg/kg/day p.o. to healthy mice during 3 consecutive days. Mice were monitored daily over a period of 30 days for survival and appearance of symptoms indicating toxicity such as ruffled fur, lethargy, loss of appetite, lacrimation, salivation, diarrhoea, convulsions and weight loss. Compounds not causing signs of gross toxicity were included in further investigations.

The in vivo antimalarial activity of the non-toxic compounds was evaluated by a modified Thompson test [14]. Mice were infected i.p. with 10⁶ infected erythrocytes in phosphate-buffered saline (PBS) (250 μ L total volume) obtained from the peripheral blood of a *P. berghei* ANKA-infected donor mouse; this time point was considered Day 0. Mice were treated with investigational compounds once a day for 3 consecutive days on Days 3, 4 and 5 post-infection. All compounds were administered p.o. at a dose of 160 mg/kg/day (200 μ L total volume) in accordance with previous experience of both our and other groups [5,15]. Infected mice treated with the same dose of CQ served as a positive control group, whereas untreated infected mice served as a negative control group. Survival and parasitaemia were monitored for 30 days post-infection.

Parasitaemia was monitored by microscopic examination of Giemsa-stained peripheral blood smears using mouse tail blood twice a week, starting immediately before the initiation of treatment (Day 3 post-infection). Up to 1000 erythrocytes were counted and parasitaemia was calculated. The sample obtained immediately before treatment initiation was of particular importance. Blood smears positive for *P. berghei* parasites presented evidence that mice were infected to a degree that would allow microscopic monitoring of treatment effects. In the absence of

Table 1

Compounds grouped according to modifications at the aminoquinoline moiety.

Compound group/compound	Compound	Reference	Compound no.
4-Aminoquinolines			
AQ4	N ¹ -(4-(5-fluorobenzo[b]thiophen-3-yl)benzyl)-N ³ -(quinolin-4-yl)propane-1,3-diamine	[11]	23
AQ5	4-(5-(4-((methyl(8-(quinolin-4-ylamino)octyl)amino)methyl)phenyl)thiophen-2-yl)benzimidazole	[11]	46
7-Chloro-4-aminoquinolines			
CIAQ7	N ¹ -(benzo[b]thiophen-3-ylmethyl)-N ² -(7-chloroquinolin-4-yl)ethane-1,2-diamine	[10]	55
CIAQ9	N ⁴ -(7-chloroquinolin-4-yl)-N ¹ -(4-(5-fluorobenzo[b]thiophen-3-yl)benzyl)pentane-1,4-diamine	[11]	25
CIAQ10	3-(((3-(7-chloroquinolin-4-yl)amino)propyl)amino)methyl)benzo[b]thiophene-6-carbonitrile	[10]	61
CIAQ11	4-(5-(4-(((8-((7-chloroquinolin-4-yl)amino)octyl)amino)methyl)phenyl)thiophen-2-yl)benzimidazole	[11]	42
CIAQ12	N ¹ -benzyl-N ² -(7-chloroquinolin-4-yl)ethane-1,2-diamine	[9]	6
CIAQ13	N ¹ -benzyl-N ³ -(7-chloroquinolin-4-yl)propane-1,3-diamine	[9]	7
CIAQ14	N ¹ -(4-bromobenzyl)-N ² -(7-chloroquinolin-4-yl)ethane-1,2-diamine	[9]	14
CIAQ15	N ¹ -(4-bromobenzyl)-N ³ -(7-chloroquinolin-4-yl)propane-1,3-diamine	[9]	15

microscopic evidence of parasites, the animal was excluded from the experiment.

Compound efficacy was evaluated on the basis of survival of treated versus untreated mice as assessed both by the rate and length of survival (expressed in % and days, respectively) and the presence and level of parasitaemia over time. Cure was defined as survival past Day 31 p.i. and clearance of parasitaemia.

2.6. Statistical analysis

IC₅₀ values were obtained using a sigmoidal dose–response model with the variable slope fitted to the results. The rates of survival in particular groups were estimated by the Kaplan–Meier product limit method and were compared by the log-rank (two curves) and log-rank for trends (three or more curves) tests. The level of statistical significance was 0.05. All statistics was performed using GraphPad Prism (GraphPad Software Inc., La Jolla, CA, USA).

3. Results

A series of ten novel aminoquinolines with modifications at the linker and at the quinoline nucleus was examined both in vitro and in vivo for antimalarial efficacy. At a concentration of 500 nM, eight (AQ4, AQ5, CIAQ9–CIAQ13 and CIAQ15) of the ten compounds inhibited the growth both of the CQ^S and CQ^R *P. falciparum* strain by ≥50%, whilst one (CIAQ7) exerted this effect only against the CQ^R strain (Fig. 1). When titrated to determine IC₅₀ values, all nine compounds were more active than CQ against the CQ^R strain, whilst none performed better than CQ against the CQ^S strain (Table 2). Although all of these nine compounds qualified for further examination, four (AQ4, AQ5, CIAQ10 and CIAQ12) were unavailable due to technical reasons (not synthesised in sufficient quantity).

The five available compounds (CIAQ7, CIAQ9, CIAQ11, CIAQ13 and CIAQ15) were first assayed for toxicity. No gross toxicity was noted in any mice given any of the compounds at a dose of 160 mg/kg/day, and all five compounds were thus subjected to in vivo testing. Prior to treatment administration, all mice were checked for parasitaemia; the pre-treatment parasitaemia was 0.7 ± 0.4% in mice assigned to the negative control group and 0.6 ± 0.3% in those assigned to treatment ($P > 0.05$). The results showed that compared with untreated control animals that died after 7.3 ± 0.5 days, treatment with all five investigational compounds significantly ($P < 0.05$) prolonged the length of survival (Table 2) to a mean of 17.7 ± 5.3 days for the 23 mice that died during the experiment. However, comparative analysis of the efficacy of the compounds showed different levels of activity (Fig. 2). Compounds CIAQ9 and CIAQ13 afforded survival of one mouse each (out of four

and five, respectively) past Day 31, but with high parasitaemia as observed on blood smears, whilst the mice that died survived 17.3 ± 0.6 days and 14.75 ± 1.7 days, respectively. On the other hand, compounds CIAQ7, CIAQ11 and CIAQ15 did not afford survival of any mice but prolonged length of survival of treated animals to 18.6 ± 8.05, 18.6 ± 7.6 and 18.3 ± 3.4 days, respectively (Table 2).

Regular monitoring of parasitaemia in each animal throughout the entire experiment allowed for several interesting observations (Tables 2 and 3). In untreated control mice at the time of death, parasitaemia ranged between 11.8% and 17.3% (mean 14.8 ± 2.3%). In contrast, the mean parasitaemia of the treated animals that died by Day 17 ($n = 16$; mean survival 14.8 ± 2.5 days) was only 4.8 ± 3.5%. Another two animals (both treated with CIAQ15) survived to Day 21 and Day 24, respectively, with a mean parasitaemia of 7.8 ± 3.3%, also significantly ($P < 0.05$) below that of the control mice. On the other hand, all other animals lived longer (five dying between Days 21 and 28, with a mean survival of 25 ± 4 days, and two still alive at Day 31), with a mean parasitaemia of 70 ± 10.5%.

4. Discussion

The results presented here demonstrate a significant antimalarial potential of the examined aminoquinoline series. Nine of the ten compounds were more active than CQ against the CQ^R *P. falciparum* strain Dd2. Moreover, a significant biological activity of five compounds (CIAQ7, CIAQ9, CIAQ11, CIAQ13 and CIAQ15) against *P. berghei* was shown. Whereas all five compounds significantly prolonged the length of survival compared with untreated infected animals, two compounds afforded survival up to Day 31 of 1/4 (CIAQ9) and 1/5 (CIAQ13) treated mice, albeit with residual parasitaemia.

All five examined compounds that exhibited significant in vivo activity share the same AQ nucleus with a chlorine atom at the C(7) position. It is known that the AQ ring is required for complexation with ferriprotoporphyrin but is not sufficient for the inhibition of haemozoin synthesis [16]. A chlorine atom at position C(7) is introduced into the AQ molecule for this purpose [16–19], and the present results confirm its role. Benzothioephene CIAQ9 is one of the two compounds that afforded survival (if only of one mouse) past Day 31. In our recent investigations of another series of novel aminoquinolines, two benzothioephenes were among the most effective compounds [8], whilst another afforded cure [11]. Taken together, these results illustrate the significance of the benzothioephene structure introduced in the side chain of aminoquinolines. Benzothioephene is suitable for the synthesis of large bioactive molecules owing to its heterocyclic structure, whilst the wide range of pharmacological activities (over 40) of this ring





Fig. 1. In vitro activity of ten novel compounds against two *Plasmodium falciparum* strains, chloroquine-sensitive *P. falciparum* strain 3D7 and chloroquine-resistant *P. falciparum* strain Dd2. Three replicates per compound were performed for each *P. falciparum* strain. LDH, lactate dehydrogenase.

Table 2

Antimalarial effect of experimental aminoquinolines examined in vitro and in vivo.

Compound group/compound	In vitro (LDH assay)		In vivo (Thompson test), treatment with 160 mg/kg/day orally during 3 days (Days 3, 4 and 5 p.i.)		
	GM IC ₅₀ (nM)		No. of mice dead on day (d) p.i.	Mice alive on Day 31/total	Treatment effect
	3D7 ^a	Dd2 ^b			
4-Aminoquinolines					
AQ4	18.77	73.35	N/A		
AQ5	119.58	71.52	N/A		
7-Chloro-4-aminoquinolines					
CIAQ7	>500	14.43	2 D12, 1 D15, 1 D24, 1 D30	0/5	Prolonged time to death ^c
CIAQ9	82.35	186.79	2 D17, 1 D18	1/4	Prolonged time to death ^c (3/4); survival ^d (1/4)
CIAQ10	50.23	18.57	N/A		
CIAQ11	97.26	174.12	1 D9, 1 D13, 1 D20, 1 D23, 1 D28	0/5	Prolonged time to death ^c
CIAQ12	32.2	40.82	N/A		
CIAQ13	22.05	37.97	1 D13, 1 D14, 1 D15, 1 D17	1/5	Prolonged time to death ^c (4/5); survival ^d (1/5)
CIAQ15	30.11	18.81	1 D15, 1 D16, 2 D17, 1 D21, 1 D24	0/6	Prolonged time to death ^c
Chloroquine (control)	18.63	270.57	0	5/5	Cure

LDH, lactate dehydrogenase; GM, geometric mean; IC₅₀, 50% inhibitory concentration; p.i., post-infection; d, day; N/A, not available.^a Chloroquine-sensitive *Plasmodium falciparum* strain 3D7.^b Chloroquine-resistant *P. falciparum* strain Dd2.^c Versus infected untreated (control) mice that died 7.33 ± 0.48 days p.i.^d With residual parasitaemia.**Fig. 2.** Effect of a 3-day oral treatment with 160 mg/kg/day of chloroquine (CQ) and five investigational compounds (CIAQ7, CIAQ9, CIAQ11, CIAQ13 and CIAQ15) on the survival of mice ($n = 5$ per compound, except for CIAQ9 and CIAQ15 where there were 4 and 6 mice, respectively) infected with *Plasmodium berghei* ANKA strain. The blue box indicates treatment days.

has been associated with the presence of a sulfur atom [20]. Among the many benzothiofene activities [21], in vitro antimalarial activity has recently been shown for benzothiofene-2-carboxamide [22] and bromobenzothiofene carboxamide derivatives [23], which were also active in vivo.

The observation of the different levels of parasitaemia associated with early or late death (and occasional survival with parasitaemia in the range from 70–90% defined as

hyperparasitaemia [24]), as the end result of treatment with compounds CIAQ7, CIAQ11 and CIAQ9, respectively (Table 3), deserves consideration. The C57BL/6 mouse strain is highly susceptible to the neurological syndrome with an incidence of cerebral malaria ranging from 60–100%, and mice die with low parasitaemia (on average < 15%) within 6–10 days after being challenged with 10^6 *P. berghei* ANKA blood-stage parasites [24]. On the other hand, in cerebral malaria-resistant mouse strains,

Table 3
Parasitaemia in individual *Plasmodium berghei*-infected mice treated orally with five experimental compounds (160 mg/kg/day) and in positive (CQ) and negative (untreated infected) control groups at various sampling points.

Compound	Parasitaemia (%)								Mouse
	Before treatment	Day 7	Day 10	Day 14	Day 17	Day 21	Day 24	Day 28	
CIAQ7	0.4	0.2	5.3	Death d12					1
	0.4	0.3	2.9	4	4.6	57	Death d24		2
	0.2	0.2	0.3	6.9	Death d15				3
	1	0.2	2.7	5	9.3	55.2	82	86	4
	0.7	0.2	7	Death d12					5
CIAQ9	0.4	0.4	0.5	0.7	Death d17				1
	0.6	0.2	0.3	0.5	1.5	2.7	52	72	2
	0.5	0.3	0.4	0.5	Death d17				3
	0.5	0.4	0.6	0.8	3.7	Death d18			4
CIAQ11	0.4	6.6	13.5	47	60	Death d20			1
	0.3	0.2	9.2	Death d13					2
	0.2	9	Death d9						3
	0.6	10	15	23.4	53.2	70	Death d23		4
	0.4	8.9	12	12.8	50	64	75	Death d28	5
CIAQ13	0.3	0	0.1	n.s.	5.1	17.1	n.s.	n.s.	1
	1.1	0	1.1	n.s.	Death d15				2
	n.s.	n.s.	n.s.	n.s.	Death d17				3
	n.s.	n.s.	n.s.	n.s.	Death d13				4
	n.s.	n.s.	n.s.	n.s.	Death d14				5
CIAQ15	1.1	0	0.1	n.s.	0.5	10.1	Death d24		1
	1.6	0	1.2	n.s.	5.5	Death d21			2
	n.s.	n.s.	n.s.	n.s.	Death d15				3
	n.s.	n.s.	n.s.	n.s.	Death d16				4
	n.s.	n.s.	n.s.	n.s.	Death d17				5
	n.s.	n.s.	n.s.	n.s.	Death d17				6
CQ	0.6	0	0	0	0	0	0	0	1
	0.8	0	0	0	0	0	0	0	2
	0.4	0	0	0	0	0	0	0	3
	0.9	0	0	0	0	0	0	0	4
	0.7	0	0	0	0	0	0	0	5
Untreated infected mice	1	17.3, Death d7							1
	0.5	Death d7							2
	0.9	Death d7							3
	0.4	15.1	Death d8					4	
	0.3	15.2	Death d8					5	
	0.2	Death d7							6
	1.2	Death d7							7
	1.2	11.8	Death d8					8	
	0.9	Death d7							9

d, day; n.s., not sampled.

death generally occurs during the third or fourth week post-infection due to anaemia caused by hyperparasitaemia [24]. However, delayed death associated with hyperparasitaemia in *P. berghei*-infected mice owing to pharmacological treatment or genetic background has been observed [25,26]. In our study, untreated infected animals died on average after 7.3 days, with a parasitaemia of 14.8%, exactly in line with the study by Bagot et al. [24]. In contrast, the investigational compounds all significantly prolonged survival, with a mean survival of mice that died during the experiment of 17.7 days. Among the mice treated with all five compounds, despite high variability within particular treatment groups, the animals clearly segregated into roughly a two-third majority that died within the first 17 days (mean 14.8 days) with a low parasitaemia of 5%, and one-third that survived a mean of 25 days with a mean parasitaemia of 70% (Fig. 3). It stands to reason that the first developed cerebral malaria as expected in *P. berghei*-infected C57BL/6 mice, but with a delay, whereas those that died at later times succumbed to anaemia caused by hyperparasitaemia. Hence, the results presented herein indicate that the compounds afforded prevention of cerebral malaria or a delay




Fig. 3. Example of peripheral blood smear at Day 24 post-infection showing hyperparasitaemia in *Plasmodium berghei*-infected C57BL/6 mouse treated with CIAQ11 (Giemsa staining, light microscope, magnification 1000 \times).

in its development in a mouse strain susceptible to cerebral malaria.

One limitation of this study is that we have not studied the residual parasite populations in mice that survived to the end of the experiment with a hyperparasitaemia of up to 86%, which could have shed more light on their virulence and pathogenicity. However, this was not possible since microscopic examination of the blood smears was not performed immediately following the sampling because of the large number of blood smears, and the residual parasites were not preserved.

In conclusion, this series of aminoquinolines exhibited significant antimalarial activity. Excellent in vitro activity against the CQ^R strain of even nine of the ten compounds is highly relevant, since the primary goal of new treatment options is to overcome parasite resistance. Interestingly, several compounds conferred resistance to cerebral malaria to mice prone to the neurological syndrome, an observation that merits further investigation.

Funding

This work was supported by grants nos. III 41019 and ON172008 from the Serbian Ministry of Education, Science and Technological Development.

Competing interests

None declared.

Ethical approval

This study was carried out in accordance with the ARRIVE guidelines and was approved by the Veterinary Directorate of the Ministry of Agriculture and Environmental Protection of Serbia [decision no. 323-07-02444/2014-05/1].

References

- [1] World Health Organization (WHO). World malaria report 2019. Geneva, Switzerland: WHO; 2019 <https://www.who.int/malaria/publications/world-malaria-report-2019/en> [accessed 26.12.19].
- [2] Ridley RG, Hofheinz W, Matile H, Jaquet C, Dorn A, Masciadri R, et al. 4-Aminoquinoline analogs of chloroquine with shortened side chains retain activity against chloroquine-resistant *Plasmodium falciparum*. *Antimicrob Agents Chemother* 1996;40:1846–54.
- [3] Madrid PB, Sherrill J, Liou AP, Weisman JL, Derisi JL, Guy RK. Synthesis of ring-substituted 4-aminoquinolines and evaluation of their antimalarial activities. *Bioorg Med Chem Lett* 2005;15:1015–8.
- [4] Šolaja BA, Opsenica D, Smith KS, Milhous WK, Terzić N, Opsenica I, et al. Novel 4-aminoquinolines active against chloroquine-resistant and sensitive *P. falciparum* strains that also inhibit botulinum serotype A. *J Med Chem* 2008;51:4388–91.
- [5] Opsenica I, Burnett JC, Gussio R, Opsenica D, Todorović N, Lanteri CA, et al. A chemotype that inhibits three unrelated pathogenic targets: the botulinum neurotoxin serotype A light chain, *P. falciparum* malaria, and the Ebola filovirus. *J Med Chem* 2011;54:1157–69.
- [6] Sinha M, Dola VR, Agarwal P, Srivastava K, Haq W, Puri SK, et al. Antiplasmodial activity of new 4-aminoquinoline derivatives against chloroquine resistant strain. *Bioorg Med Chem Lett* 2014;22:3573–86.
- [7] Kondaparla S, Soni A, Manhas A, Srivastava K, Puri SK, Katti SB. Antimalarial activity of novel 4-aminoquinolines active against drug resistant strains. *Bioorg Chem* 2016;70:74–85.
- [8] Srbljanović J, Štajner T, Konstantinović J, Terzić-Jovanović N, Uzelac A, Bobić B, et al. Examination of the antimalarial potential of experimental aminoquinolines: poor in vitro effect does not preclude in vivo efficacy. *Int J Antimicrob Agents* 2017;50:461–6.
- [9] Opsenica IM, Tot M, Gomba L, Nuss JE, Sciotti RJ, Bavari S, et al. 4-Amino-7-chloroquinolines: probing ligand efficiency provides botulinum neurotoxin serotype A light chain inhibitors with significant antiprotozoal activity. *J Med Chem* 2013;56:5860–71.
- [10] Terzić N, Konstantinović J, Tot M, Burojević J, Djurković-Djaković O, Srbljanović J, et al. Re-investigation of old pharmacophores: are aminoquinolines and tetraoxanes potential two-stage antimalarials? *J Med Chem* 2016;59:264–81.
- [11] Konstantinović J, Videnović M, Srbljanović J, Djurković-Djaković O, Bogojević K, Sciotti R, et al. Antimalarials with benzothioephene moieties as aminoquinoline partners. *Molecules* 2017;22:343, doi:<http://dx.doi.org/10.3390/molecules.22030343>.
- [12] Trager W, Jensen JB. Human malaria parasites in continuous culture. *Science* 1976;193:673–5.
- [13] Makler MT, Ries JM, Williams JA, Bancroft JE, Piper RC, Gibbins BL, et al. Parasite lactate dehydrogenase as an assay for *Plasmodium falciparum* drug sensitivity. *Am J Trop Med Hyg* 1993;48:739–41.
- [14] Ager AL. Experimental models: rodent malaria models (in vivo). In: Peters W, Richards WHG, editors. *Handbook of experimental pharmacology: antimalarial drugs*. New York, NY: Springer Verlag; 1983. p. 225–54.
- [15] Sáenz FE, Mutka T, Udenze K, Oduola AMJ, Kyle DE. Novel 4-aminoquinoline analogs highly active against the blood and sexual stages of *Plasmodium* in vivo and in vitro. *Antimicrob Agents Chemother* 2012;56:4685–92.
- [16] Egan TJ, Hunter R, Kaschula CH, Marques HM, Misplon A, Walden J. Structure–function relationships in aminoquinolines: effect of amino and chloro groups on quinoline–hematin complex formation, inhibition of β -hematin formation, and antiplasmodial activity. *J Med Chem* 2000;43:283–91.
- [17] De D, Krogstad FM, Byers LD, Krogstad DJ. Structure–activity relationships for antiplasmodial activity among 7-substituted 4-aminoquinolines. *J Med Chem* 1998;41:4918–26.
- [18] Kaschula CH, Egan TJ, Hunter R, Basilico N, Parapini S, Taramelli D, et al. Structure–activity relationships in 4-aminoquinoline antiplasmodials. The role of the group at the 7-position. *J Med Chem* 2002;45:3531–9.
- [19] Hwang JY, Kawasuji T, Lowes DJ, Clark JA, Connelly MC, Zhu F, et al. Synthesis and evaluation of 7-substituted 4-aminoquinoline analogs for antimalarial activity. *J Med Chem* 2011;54:7084–93.
- [20] Kharb R, Bansal K. Perspectives on antimicrobial potential of benzothioephene derivatives. *Res J Pharm Biol Chem Sci* 2013;4:1585–606.
- [21] Keri RS, Chand K, Budagumpi S, Balappa Somappa S, Patil SA, Nagaraja BM. An overview of benzo[b]thiophene-based medicinal chemistry. *Eur J Med Chem* 2017;138:1002–33.
- [22] Pieroni M, Azzali E, Basilico N, Parapini S, Zolkiewski M, Beato C, et al. Accepting the invitation to open innovation in malaria drug discovery: synthesis, biological evaluation, and investigation on the structure–activity relationships of benzo[b]thiophene-2-carboxamides as antimalarial agents. *J Med Chem* 2017;60:1959–70.
- [23] Banerjee T, Kapoor N, Surolia N, Surolia A. Benzothioephene carboxamide derivatives as novel antimalarials. *IUBMB Life* 2011;63:1111–5.
- [24] Bagot S, Boubou MI, Campino S, Behrschmidt C, Gorgette O, Guénet JL, et al. Susceptibility to experimental cerebral malaria induced by *Plasmodium berghei* ANKA in inbred mouse strains recently derived from wild stock. *Infect Immun* 2002;70:2049–56.
- [25] Blavid R, Kusch P, Hauber J, Eschweiler U, Sarite SR, Specht S, et al. Down-regulation of hypusine biosynthesis in *Plasmodium* by inhibition of S-adenosyl-methionine-decarboxylase. *Amino Acids* 2010;38:461–9.
- [26] Campino S, Bagot S, Bergman ML, Almeida P, Sepúlveda N, Pied S, et al. Genetic control of parasite clearance leads to resistance to *Plasmodium berghei* ANKA infection and confers immunity. *Genes Immun* 2005;6:416–21.